



Changes in gut microbiota control inflammation in obese mice through a mechanism involving GLP-2-driven improvement of gut permeability

P D Cani, S Possemiers, T Van de Wiele, Y Guiot, A Everard, O Rottier, L Geurts, D Naslain, A Neyrinck, D M Lambert, G G Muccioli and N M Delzenne

Gut 2009;58;1091-1103; originally published online 24 Feb 2009;
doi:10.1136/gut.2008.165886

Updated information and services can be found at:
<http://gut.bmj.com/cgi/content/full/58/8/1091>

	<i>These include:</i>
Data supplement	"Web only appendices" http://gut.bmj.com/cgi/content/full/gut.2008.165886/DC1
References	This article cites 84 articles, 33 of which can be accessed free at: http://gut.bmj.com/cgi/content/full/58/8/1091#BIBL 1 online articles that cite this article can be accessed at: http://gut.bmj.com/cgi/content/full/58/8/1091#otherarticles
Open Access	This article is free to access
Email alerting service	Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Topic collections Articles on similar topics can be found in the following collections
[Unlocked \(151 articles\)](#)

Notes

To order reprints of this article go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to *Gut* go to:
<http://journals.bmj.com/subscriptions/>

